

THE PROMISE

OF *illumigene*[®]

Group B *Streptococcus* (GBS)

Empower your lab with the speed and accuracy of molecular testing for Group B Streptococcus

The Concern:

- Group B *Streptococcus* is a leading cause of illness and death among newborns¹
- 61.4% of term infants with GBS disease were born to women who had tested negative for Group B *Streptococcus* before delivery²
- Sensitivity of conventional culture is as low as 42.3%³
- Conventional culture can take 3-4 days to complete⁴

The Solution:

- *illumigene*[®] GBS provides definitive and accurate results for improved patient management
- *illumigene*[®] GBS allows labs to fully comply with the 2010 CDC guidelines
- Broth enrichment followed by *illumigene*[®] GBS provides increased sensitivity over culture for the detection of GBS⁵ —48/51 culture negative specimens were positive by another commercially available molecular method



SENSITIVITY	SPECIFICITY
97.4%	92.3%


ORDER TODAY



TEST TOMORROW

 **Meridian**
Bioscience, Inc.
Inspired Science. Trusted Solutions.[®]

illumigene® GBS Assay Procedure



Ordering information

<i>illumigene</i> ® Group B Streptococcus Assay.....	280350
<i>illumigene</i> ® Group B Streptococcus External Control Kit.....	279900



For more information, contact an *illumigene*® specialist at 1-888-763-6769 or visit us on the web at www.meridianbioscience.com.

REFERENCES

- Centers for Disease Control and Prevention. Prevention of Perinatal Group B *Streptococcal* Disease. MMWR 2010;59(No. RR-10):pp. 1-32.
- M. Van Dyke, Ph.D., et al. 2009. Evaluation of Universal Antenatal Screening for Group B *Streptococcus*. The New England Journal of Medicine. 360:pp. 2626-2636.
- F. Rallu, P. Barraja, C. Scrivo, V. Martel-Laferriere, and C. Laferriere. 2006. Sensitivities of Antigen Detection and PCR assays Greatly Increased Compared to that of the Standard Culture Method for Screening for Group B *Streptococcus* Carriage in Pregnant Women. Journal of Clinical Microbiology. 44: pp. 725-728.
- N. Montague, T. Cleary, O. Martinez, and G. Procop. 2008. Detection of Group B *Streptococci* in Lim Broth by Use of Group B *Streptococcus* Peptide Nucleic Acid Fluorescent in Situ Hybridization and Selective and Nonselective Agars. Journal of Clinical Microbiology. 46: pp. 3470-3472.
- illumigene*® GBS package insert, July 2011.

